

Vaccine-Specific Coverage in NC among Children 19-35 months, National Immunization Survey, 2011-2013

Estimated Vaccination Coverage NIS (19 through 35 months)									
	2011			2012			2013		
Vaccine(s)	US %	NC %	NC Rank	US %	NC %	NC Rank	US %	NC %	NC Rank
DTaP (4 doses)	84.6	81.3	39	82.5	85.9	15	83.1	87.5	12
Polio (3 doses)	93.9	93.5	31	92.8	92.8	33	92.7	95.9	9
MMR (1 dose)	91.6	92.3	21	90.8	89.0	41	91.9	96.0	3
Hib*	94.2	93.0	34	80.9	83.0	18	82.0	87.0	7
Hepatitis B (birth dose)	68.6	75.0	16	71.6	78.2	17	74.2	82.1	6
Hepatitis B (3 doses)	91.1	90.8	27	89.7	89.2	32	90.8	94.3	8
Var (1 dose)	90.8	89.6	28	90.2	88.0	38	91.2	94.6	5
PCV (4 doses)^	84.4	84.5	25	81.9	82.0	29	82.0	86.5	15
Hepatitis A (2 doses)^	52.2	40.8	48	53.0	48.5	38	54.7	51.6	34
Rotavirus (complete)^	67.3	70.5	16	68.6	68.0	26	72.6	75.4	15
Overall 4:3:1:3:3:1#	77.0	75.3	34	71.9	76.2	11	74.0	76.6	13
Overall 4:3:1:3:3:1:4†	73.3	73.3	24	68.4	75.4	5	70.4	72.0	20

*Due to a nationwide shortage, chart refers to primary Hib series (2 or 3 doses, depending on the product used) in 2011. 2012 and 2013 estimates refer to the full Hib series (3 or 4 doses, depending on the product used).

^Hepatitis A and Rotavirus vaccines are not required in North Carolina. Pneumococcal was not required during this time frame.

#The measured 4:3:1:3:3:1 series was at least 4 DTaP, 3 Polio, 1 MMR, 3 Hib, 3 Hep B, 1 Var. It is referred to as 4:3:1:4:3:1 series in 2013 NIS data.

†The measured 4:3:1:3:3:1:4 series was the above series plus at least 4 PCV. It is referred to as 4:3:1:4:3:1:4 series in 2013 NIS data.