

**NORTH CAROLINA IMMUNIZATION PROGRAM (NCIP)
COVERAGE CRITERIA FOR ALL CHILDREN PRESENT IN NORTH CAROLINA**

Effective: November 17, 2017

The purpose of this document is to distinguish which cohorts of persons, present in North Carolina, are eligible for vaccine from the North Carolina Immunization Program (NCIP). NCIP vaccine is generally available for children through the age of 18 in compliance with recommendations made by the Advisory Committee on Immunization Practices (ACIP) and the Vaccines for Children (VFC) program. Exceptions for adult vaccine usages are noted below. ACIP recommends certain vaccines for certain cohorts who are not covered by NCIP vaccine. Health care providers must use privately purchased vaccine for those cohorts who wish to have vaccine and are not covered by this coverage criteria. Questions about current recommendations for each vaccine can be found at: <https://www.cdc.gov/vaccines/hcp/acip-recs/index.html>. Local Health Departments (LHD); Federally Qualified Health Centers (FQHC) and Rural Health Clinics (RHC) are noted throughout when there is vaccine usage specific to their facilities.

VACCINE	AGES COVERED	ELIGIBILITY	CRITERIA FOR NCIP VACCINE USAGE
DT Pediatric	≥2 months through 6 years	UNIVERSAL	DT Pediatric can be ordered for any child, ≥2 months through 6 years, that has a known medical contraindication to any pertussis containing vaccine.
DTaP	≥2 months through 6 years	VFC ONLY	
DTaP, Hep B, and polio combination (Pediatrix®)	≥2 months through 6 years	VFC ONLY	
DTaP, polio, and Hib combination (Pentacel®)	≥2 months through 4 years	VFC ONLY	
DTaP/IPV combination (Kinrix®; Quadracel®)	≥4 years through 6 years	VFC ONLY	Only to be used for children ≥ 4 years through 6 years of age. Kinrix as the 5 th dose in the DTaP series, & as the 4 th in the IPV series. Quadracel as the 5 th dose in the DTaP series, & as the 4 th or 5 th dose in the IPV series
Hepatitis A Pediatric use	12 months through 18 years	VFC ONLY	
Hepatitis B Pediatric use	Birth through 18 years	VFC ONLY	<ol style="list-style-type: none"> 1. The birth dose of Hepatitis B vaccine is available universally. 2. Other doses of hepatitis B vaccine may be given to children through 18 years of age who are VFC-eligible only. 3. LHD Only: Children through 18 years of age, regardless of insurance status who are household, sexual or needle sharing contacts of an acute or chronic Hepatitis B infected person may receive Hepatitis B vaccine at the LHD. All household, sexual, or needle sharing contacts of an acute or chronic Hepatitis B infected person must be referred to the LHD for testing and vaccination.
Hepatitis B Pediatric Product (Engerix-B®) (Recombivax®)	Certain adults 19 years of age	UNINSURED 19 YEARS OF AGE	<ol style="list-style-type: none"> 1. Any uninsured adult 19 years of age may receive state-supplied Hep B if the first dose of Hep B is administered before the 19th birthday. The series must be complete before the 20th birthday if using state-supplied vaccine. 2. LHD Only: Uninsured immigrants with refugee status who are from endemic countries who are 19 years of age may receive Hepatitis B vaccine at the LHD. All immigrants with refugee status must be referred to the LHD for testing and vaccination. 3. LHD Only: Adults 19 years of age, regardless of insurance status, who are household, sexual or needle sharing contacts of an acute or chronic Hepatitis B infected person may receive Hepatitis B vaccine at the LHD. All household, sexual, or needle sharing contacts of an acute or chronic Hepatitis B infected person must be referred to the LHD for testing/vaccination.

**NORTH CAROLINA IMMUNIZATION PROGRAM (NCIP)
COVERAGE CRITERIA FOR ALL CHILDREN PRESENT IN NORTH CAROLINA**

Effective: November 17, 2017

VACCINE	AGES COVERED	ELIGIBILITY	CRITERIA FOR NCIP VACCINE USAGE
Hepatitis B Adult use	Certain adults \geq 20 years may receive state-supplied Hep B	ADULT USE	LHD Only: Adult state-supplied Hepatitis B is available for: <ol style="list-style-type: none"> Individuals who are \geq20 years of age, regardless of insurance status, who are household, sexual or needle sharing contacts of an acute or chronic Hepatitis B infected person may receive Hepatitis B vaccine at the LHD. All household, sexual, or needle sharing contacts of an acute or chronic Hepatitis B infected person must be referred to the LHD for testing and vaccination. Uninsured immigrants with refugee status who are from endemic countries who are \geq 20 years of age may receive Hepatitis B vaccine at the LHD. All immigrants with refugee status must be referred to the LHD for testing and vaccination.
HepA/HepB Combination (Twinrix®)	\geq 18 years	UNINSURED ADULT USE	<u>LHD/FQHC/RHC Only:</u> Any uninsured adult who meets one or more of the ACIP recommended coverage groups can receive a three-dose series of the combination Hep A/Hep B vaccine at the LHD, FQHC, or RHC. State-supplied Hep A/Hep B vaccine cannot be used for the accelerated schedule, four dose series or for persons with a documented history of a completed hepatitis A or B series.
Hib	\geq 2 months through 4 years	VFC ONLY	Certain high-risk children >59 months through 18 years of age, who are not appropriately vaccinated, may receive one dose.
9vHPV (Gardasil® 9)	Females and males 9 through 18 years	VFC ONLY	State-supplied Gardasil® (9-valent) vaccines for HPV are available for males and females 9 through 18 years of age. The series must be completed prior to the 19th birthday.
Influenza Pediatric use	>6 months through 18 years	VFC ONLY	
Influenza Adult use IIV4 (Fluzone® 0.5 mL pre-filled syringes only)	Certain adults, (19 and older) without insurance	UNINSURED FEMALE ADULT USE	<u>LHD/FQHC/RHC Only:</u> State-supplied influenza vaccine is available for: <ol style="list-style-type: none"> Non-Medicaid, uninsured women who are pregnant during flu season, and receiving services at the LHD, FQHC, or RHC. Females enrolled in the Be Smart family planning program receiving services at the LHD, FQHC, or RHC.
Meningococcal Conjugate (Menacta®) (Menveo®)	2 months through 10 years	VFC ONLY	Available for children with high-risk conditions. Menactra® is licensed starting at 9 months of age. Menveo® starts at 2 months of age.
Meningococcal Conjugate (Menacta®) (Menveo®)	11 through 18 years	VFC ONLY	
Meningococcal Conjugate Adult use	Adults 19 years through 22 years	UNINSURED ADULT USE	A single dose of meningococcal vaccine is available for uninsured unvaccinated first year college students <u>living in dormitories.</u>
Meningococcal Serogroup B Bexsero® Trumenba®	10 through 18 years	VFC ONLY	MenB vaccines are recommended & available for children \geq 10 through 18 years of age at increased risk of disease. MenB may be administered to adolescents aged 16–18 years of age who are not at increased risk for meningococcal disease based on the provider’s clinical discretion. Follow ACIP guidance.
MMR Pediatric use	\geq 12 months through 18 years	VFC ONLY	

**NORTH CAROLINA IMMUNIZATION PROGRAM (NCIP)
COVERAGE CRITERIA FOR ALL CHILDREN PRESENT IN NORTH CAROLINA**

Effective: November 17, 2017

VACCINE	AGES COVERED	ELIGIBILITY	ELIGIBILITY CRITERIA FOR NCIP VACCINE USAGE
MMR Adult use	Certain adults (19 and older) without insurance	UNINSURED ADULT USE	<p>One dose of MMR vaccine is available for:</p> <ol style="list-style-type: none"> LHD only: Uninsured immigrants with refugee status of any age without documentation of a prior MMR may receive MMR at the LHD. LHD/FQHC/RHC only: All uninsured susceptible persons without documentation of a prior MMR may receive MMR at the LHD, FQHC, or RHC. <p>Two doses of MMR vaccine are available for:</p> <ol style="list-style-type: none"> Any uninsured person entering a four-year college or university in North Carolina for the first time, in addition to, any uninsured adults attending any other post-high school educational institutions (i.e. community colleges and technical schools because: <ul style="list-style-type: none"> ➤ Two doses of measles containing vaccine is required by law for all persons born in 1957 or later entering a North Carolina college/university*** for the first time on or after July 1, 1994 and that ➤ Two doses of mumps are also recommended and available for all uninsured adult students attending post-high school educational institutions (i.e. community colleges and technical schools). LHD only: All uninsured women ages 19-44 years at the LHD.
MMRV	≥12 months through 12 years	VFC ONLY	
IPV-Polio	≥2 months through 17 years	VFC ONLY	
Pneumococcal Conjugate 13-valent (PCV13)	≥2 through 59 months	VFC ONLY	
Pneumococcal Conjugate 13-valent (PCV13)	≥60 months through 18 years	VFC ONLY	Children ≥60 months through 18 years with certain high-risk conditions.
Pneumococcal Polysaccharide 23-valent (PPSV23)	≥2 years through 18 years	VFC ONLY	Children ≥2 years through 18 years with certain high-risk conditions.
Rotavirus	≥6 weeks through 7 months	VFC ONLY	
Td Pediatric use	≥7 years through 18 years	VFC ONLY	
Td Adult use	Certain adults (19 and older) without insurance	UNINSURED ADULT USE	<ol style="list-style-type: none"> Can be given to any uninsured person 19 years of age or older entering a North Carolina college/university*** (where Td is required by NC Immunization Law) for the first time and has not had a booster in 10 years, or has not completed the 3 dose primary series. LHD/FQHC/RHC: Any uninsured person 19 years of age or older can receive Td at the LHD, FQHC, or RHC. Can be given to any uninsured person 19 years of age or older being seen in a hospital emergency department for wound management.

**NORTH CAROLINA IMMUNIZATION PROGRAM (NCIP)
COVERAGE CRITERIA FOR ALL CHILDREN PRESENT IN NORTH CAROLINA**

Effective: November 17, 2017

VACCINE	AGES COVERED	ELIGIBILITY	ELIGIBILITY CRITERIA FOR NCIP VACCINE USAGE
Tdap Pediatric use	≥7 years through 18 years	VFC ONLY	
Tdap Adult use	Certain adults, (19 and older) without insurance	UNINSURED ADULT USE	<ol style="list-style-type: none"> One dose of state-supplied vaccine may be administered to any uninsured adult for whom Tdap is indicated. LHD only: State-supplied Tdap may administer to uninsured pregnant women 19 years and older (at any time during pregnancy regardless of prior vaccine history of Tdap, but preferably between 27-36 weeks' gestation). In addition, uninsured postpartum women 19 years and older may receive one dose of state supplied Tdap if there is no documentation of a previous Tdap vaccination. Both pregnant and postpartum women must be being served by the LHD in any capacity.
Varicella (Chickenpox)	> 12 months through 18 years	VFC ONLY	
Varicella (Chickenpox)	Certain adults, (19 and older) without insurance	UNINSURED FEMALE ADULT USE	<p>LHD ONLY: One dose of varicella vaccine is available for:</p> <ol style="list-style-type: none"> Uninsured women ages 19-44 years at the LHD.

VFC- Vaccines for Children: Children birth through 18 years of age that meet at least one of the following criteria are eligible for VFC vaccine:

- Medicaid enrolled - a child who is eligible or enrolled in the Medicaid program.
- Uninsured - a child who has no medical insurance coverage
- American Indian or Alaskan Native
- Underinsured (Can only be served by deputized providers such as LHD/FQHC/RHC). Underinsured include:
 - Children who have commercial (private) health insurance but the coverage does not include vaccines,
 - Children whose insurance covers only selected vaccines (VFC-eligible for non-covered vaccines only),
 - Children whose insurance caps vaccine coverage at a certain amount - once that coverage amount is reached, these children are categorized as underinsured

Children who are covered by North Carolina Health Choice (NCHC) are considered insured, with one exception: NCHC children who are American Indian or Alaskan Native are eligible for VFC vaccines. **Note:** Children whose health insurance covers vaccinations are not eligible for VFC vaccines, even when a claim for the cost of the vaccine and its administration would be denied for payment by the insurance carrier because the plan's deductible had not been met or because the insurance did not cover the total cost of the vaccine

VFC vaccine cannot be used for allergy skin testing prior to vaccination.

**** ACIP Recommendation Highlights** The ACIP recommendations for each vaccine are located on the CDC website at <http://www.cdc.gov/vaccines/hcp/acip-recs/index.html>. Full and final recommendations are published as MMWR Recommendations and Reports. Updates are often sent out as Provisional Recommendations with the most pertinent information included before the final recommendations are released. There is also separate documentation for **Contraindication/Precautions** for all vaccines on the CDC web site at: <https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html>.

***Students attending colleges within the North Carolina Community College System are exempt from North Carolina Immunization Law for college/university entry. However, some programs at community colleges, such as medical programs, may have requirements specific to their program, but this does not allow them to receive state supplied vaccine unless it is specifically stated in the coverage criteria.

*****LHD/FQHC/RHC only:** Persons covered by the Be Smart Family Planning Program are considered uninsured and may receive certain state-supplied vaccines as noted in this coverage criteria for uninsured adults if receiving services at a Local Health Department, Federally Qualified Health Center, or Rural Health Clinic. Unaccompanied minors (through 18 years of age) without proof of insurance presenting to a local health department Title X clinic can also be vaccinated with VFC vaccines.

Local health departments (LHD) who are considering using VFC vaccine for offsite clinics, must contact the Immunization Branch for specific guidance prior to scheduling the clinic. Proper storage and handling is critical at all times.

Unless specifically stated above, no NCIP vaccine may be administered to an insured individual unless the patient is an underinsured child at an FQHC, RHC, LHD, or deputized provider.