Children in Child Care

Children attending child care centers or homes must receive shots – or immunizations.

You must collect shot records from parents and update your records until all required shots have been given.

INPORTANT

Parents or guardians must provide a child's certificate of immunization within 30 days of attendance. Children who are behind in their shots may be enrolled, but must also prove they are trying to get the needed vaccines and are meeting scheduled appointments.

If you have questions about immunizations, call your local health department or contact the N/C Immunization Branch at 919-707-5550

or visit AAAA.immunize.ne.gov.

N.C. Department of Health and Human Services Division of Public Health Women's & Children's Health Section Immunization Branch

N.C. DHHS does not discriminate on the basis of race, color, national origin, sex, religion, age or disability in employment or the provision of services.

Revised 11/14

What Shots Do They Need?

By This Age

			~				
3 months	1 DTaP	1 Polio	1 Hib	1 PCV		1 Hep B	
5 months	2 DTaP	2 Polio	2 Hib	2 PCV		2 Hep B	
7 months	3 DTaP	2 Polio	2-3 Hib	3 PCV		2 Hep B	
12-16 months	3 DTaP	2 Polio	3-4 Hib	4 PCV	1 MMR	2 Hep B	1 Var⁺
19 months	4 DTaP	3 Polio	3-4 Hib	4 PCV	1 MMR	3 Нер В	1 Var+
4 years or older (in daycare only)	4 DTaP	3 Polio	3-4 Hib**	4 PCV**	1 MMR	3 Нер В	1 Var+
4 years or older (and in kindergarten)	5 DTaP*	4 Polio*	3-4 Hib**	4 PCV**	2 MMR*	3 Нер В	2 Var+*

Children Need These Shots

- * The 5th DTaP: 4th Polio; 2nd MMR; and 2nd Varicella are not required until children enter kindergarten. The last DTaP and Polio must be after age 4.
- ** Children beyond their 5th birthday are not required to receive Hib or PCV vaccines
- ⁺ Vaccination is required unless signed documentation of disease history is provided by a health care worker stating the date or age of child's infection.

Note: Hepatitis A, Rotavirus and Flu vaccines are not required or reportable but are recommended by the Advisory Committee on Immunization Practices.

Diphtheria, Tetanus and acellular Pertussis is sometimes combined with other vaccines and is written as DTaP-HepB-IPV, DTaP-IPV/Hib or DTaP-IPV. If it is written as DT, ask the parent for a doctor's note stating why DT was used instead of DTaP. Children who receive the fourth DTaP after age 4 do not need a fifth dose.

Polio Polio may be written as either OPV, IPV, Sabin Polio or EIPV. Polio may also be written as DTaP-HebB-IPV, DTaP-IVP/Hib or DTaP-IPV as a part of a combination vaccine. The fourth dose of polio must be received after the 4th birthday and before entering kindergarten. Children who receive the third dose of polio after age 4 do not need a fourth dose.

HID Haemophilus influenzae type B may be written by itself or as DTaP-IPV/Hib, or Hib-MenCY. Children receive either 3 or 4 doses of Hib, depending on the type of Hib given.

Hep \mathcal{F} Hepatitis B is required for all children born after July 1, 1994.

Pheumococcal Conjugate Vaccine is required for all children born on or after July 1, 2015. The number of doses required varies based on the age of the child's first vaccine.

MMK Measles, Mumps and Rubella is a 3-in-one shot. The first dose can be given no earlier than the 1st birthday. The 2nd dose must be given before the child enters kindergarten.

Varicella (Chickenpox) – The first dose can be given no earlier than the 1st birthday and beginning July 1, 2015, the second dose is required before entering kindergarten.